

**PLASMA PROTEASE C1 BIOPOLYMER
MARKERS PREDICTIVE OF ALZHEIMERS DISEASE**

FIELD OF THE INVENTION

This invention relates to the field of characterizing the existence of a disease state; particularly to the utilization of mass spectrometry to elucidate particular biopolymer markers indicative or predictive of a particular disease state, and most particularly to specific biopolymer markers whose up-regulation, down-regulation, or relative presence in disease vs. normal states has been determined to be useful in disease state assessment and therapeutic target recognition, development and validation.

BACKGROUND OF THE INVENTION

Methods utilizing mass spectrometry for the analysis of a target polypeptide have been taught wherein the polypeptide is first solubilized in an appropriate solution or reagent system. The type of solution or reagent system, e.g., comprising an organic or inorganic solvent, will depend on the properties of the polypeptide and the type of mass spectrometry performed and are well-known in the art (see, e.g., Vorm et al. (1994) Anal. Chem. 66:3281 (for MALDI) and Valaskovic et al. (1995) Anal. Chem. 67:3802 (for ESI). Mass spectrometry of peptides is further disclosed, e.g., in WO

1 93/24834 by Chait et al.

2 In one prior art embodiment, the solvent is chosen so
3 that the risk that the molecules may be decomposed by the
4 energy introduced for the vaporization process is
5 considerably reduced, or even fully excluded. This can be
6 achieved by embedding the sample in a matrix, which can be an
7 organic compound, e.g., sugar, in particular pentose or
8 hexose, but also polysaccharides such as cellulose. These
9 compounds are decomposed thermolytically into CO₂ and H₂O so
10 that no residues are formed which might lead to chemical
11 reactions. The matrix can also be an inorganic compound,
12 e.g., nitrate of ammonium which is decomposed practically
13 without leaving any residues. Use of these and other solvents
14 are further disclosed in U.S. Pat. No. 5,062,935 by Schlag et
15 al.

16 Prior art mass spectrometer formats for use in analyzing
17 the translation products include ionization (I) techniques,
18 including but not limited to matrix assisted laser desorption
19 (MALDI), continuous or pulsed electrospray (ESI) and related
20 methods (e.g., IONSPRAY or THERMOSPRAY), or massive cluster
21 impact (MCI); these ion sources can be matched with detection
22 formats including linear or non-linear reflection time-of-
23 flight (TOF), single or multiple quadropole, single or

1 multiple magnetic sector, Fourier Transform ion cyclotron
2 resonance (FTICR), ion trap, and combinations thereof (e.g.,
3 ion-trap/time-of-flight). For ionization, numerous
4 matrix/wavelength combinations (MALDI) or solvent
5 combinations (ESI) can be employed. Subattomole levels of
6 protein have been detected, for example, using ESI
7 (Valaskovic, G. A. et al., (1996) Science 273:1199-1202) or
8 MALDI (Li, L. et al., (1996) J. Am. Chem. Soc. 118:1662-1663)
9 mass spectrometry.

10 ES mass spectrometry has been introduced by Fenn et al.
11 (J. Phys. Chem. 88, 4451-59 (1984); PCT Application No. WO
12 90/14148) and current applications are summarized in recent
13 review articles (R. D. Smith et al., Anal. Chem. 62, 882-89
14 (1990) and B. Ardrey, Electrospray Mass Spectrometry,
15 Spectroscopy Europe, 4, 10-18 (1992)). MALDI-TOF mass
16 spectrometry has been introduced by Hillenkamp et al.
17 ("Matrix Assisted UV-Laser Desorption/Ionization: A New
18 Approach to Mass Spectrometry of Large Biomolecules,"
19 Biological Mass Spectrometry (Burlingame and McCloskey,
20 editors), Elsevier Science Publishers, Amsterdam, pp. 49-60,
21 1990). With ESI, the determination of molecular weights in
22 femtomole amounts of sample is very accurate due to the
23 presence of multiple ion peaks which all could be used for

1 the mass calculation.

2 The mass of the target polypeptide determined by mass
3 spectrometry is then compared to the mass of a reference
4 polypeptide of known identity. In one embodiment, the target
5 polypeptide is a polypeptide containing a number of repeated
6 amino acids directly correlated to the number of
7 trinucleotide repeats transcribed/translated from DNA; from
8 its mass alone the number of repeated trinucleotide repeats
9 in the original DNA which coded it, may be deduced.

10 U.S. Patent No. 6,020,208 utilizes a general category of
11 probe elements (i.e., sample presenting means) with Surfaces
12 Enhanced for Laser Desorption/Ionization (SELDI), within
13 which there are three (3) separate subcategories. The SELDI
14 process is directed toward a sample presenting means (i.e.,
15 probe element surface) with surface-associated (or surface-
16 bound) molecules to promote the attachment (tethering or
17 anchoring) and subsequent detachment of tethered analyte
18 molecules in a light-dependent manner, wherein the said
19 surface molecule(s) are selected from the group consisting of
20 photoactive (photolabile) molecules that participate in the
21 binding (docking, tethering, or crosslinking) of the analyte
22 molecules to the sample presenting means (by covalent
23 attachment mechanisms or otherwise).

1 analysis by a highly trained individual to determine disease
2 state versus the perception of non-disease or normal
3 physiology.

4 Richter et al, Journal of Chromatography B, 726(1999)
5 25-35, refer to a database established from human
6 hemofiltrate comprised of a mass database and a sequence
7 database. The goal of Richter et al was to analyze the
8 composition of the peptide fraction in human blood. Using
9 MALDI-TOF, over 20,000 molecular masses were detected
10 representing an estimated 5,000 different peptides. The
11 conclusion of the study was that the hemofiltrate (HF)
12 represented the peptide composition of plasma. No
13 correlation of peptides with relation to normal and/or
14 disease states is made.

15 As used herein, "analyte" refers to any atom and/or
16 molecule; including their complexes and fragment ions. The
17 term may refer to a single component or a set of components.
18 In the case of biological molecules/macromolecules or
19 "biopolymers", such analytes include but are not limited to:
20 polypeptides, polynucleotides, proteins, peptides,
21 antibodies, DNA, RNA, carbohydrates, steroids, and lipids,
22 and any detectable moiety thereof, e.g. immunologically
23 detectable fragments. Note that most important biomolecules

1 under investigation for their involvement in the structure or
2 regulation of life processes are quite large (typically
3 several thousand times larger than H₂O).

4 As used herein, the term "molecular ions" refers to
5 molecules in the charged or ionized state, typically by the
6 addition or loss of one or more protons (H⁺).

7 As used herein, the term "molecular fragmentation" or
8 "fragment ions" refers to breakdown products of analyte
9 molecules caused, for example, during laser-induced
10 desorption (especially in the absence of added matrix).

11 As used herein, the term "solid phase" refers to the
12 condition of being in the solid state, for example, on the
13 probe element surface.

14 As used herein, "gas" or "vapor phase" refers to
15 molecules in the gaseous state (i.e., in vacuo for mass
16 spectrometry).

17 As used herein, the term "analyte desorption/ionization"
18 refers to the transition of analytes from the solid phase to
19 the gas phase as ions. Note that the successful
20 desorption/ionization of large, intact molecular ions by
21 laser desorption is relatively recent (circa 1988)--the big
22 breakthrough was the chance discovery of an appropriate
23 matrix (nicotinic acid).

1 As used herein, the term "gas phase molecular ions"
2 refers to those ions that enter into the gas phase. Note that
3 large molecular mass ions such as proteins (typical
4 mass=60,000 to 70,000 times the mass of a single proton) are
5 typically not volatile (i.e., they do not normally enter into
6 the gas or vapor phase). However, in the procedure of the
7 present invention, large molecular mass ions such as proteins
8 do enter the gas or vapor phase.

9 As used herein in the case of MALDI, the term "matrix"
10 refers to any one of several small, acidic, light absorbing
11 chemicals (e.g., CHCA (alpha-cyano-4-hydroxy-cinnamic acid),
12 nicotinic or sinapinic acid) that is mixed in solution with
13 the analyte in such a manner so that, upon drying on the
14 probe element, the crystalline matrix-embedded analyte
15 molecules are successfully desorbed (by laser irradiation)
16 and ionized from the solid phase (crystals) into the gaseous
17 or vapor phase and accelerated as intact molecular ions. For
18 the MALDI process to be successful, analyte is mixed with a
19 freshly prepared solution of the chemical matrix (e.g.,
20 10,000:1 matrix:analyte) and placed on the inert probe
21 element surface to air dry just before the mass spectrometric
22 analysis. The large fold molar excess of matrix, present at
23 concentrations near saturation, facilitates crystal formation

1 and entrapment of analyte.

2 As used herein, "energy absorbing molecules (EAM)"
3 refers to any one of several small, light absorbing chemicals
4 that, when presented on the surface of a probe, facilitate
5 the neat desorption of molecules from the solid phase (i.e.,
6 surface) into the gaseous or vapor phase for subsequent
7 acceleration as intact molecular ions. The term EAM is
8 preferred, especially in reference to SELDI. Note that
9 analyte desorption by the SELDI process is defined as a
10 surface-dependent process (i.e., neat analyte may be placed
11 on a surface composed of bound EAM or EAM and analyte may be
12 mixed prior to placement on a surface). In contrast, MALDI is
13 presently thought to facilitate analyte desorption by a
14 volcanic eruption-type process that "throws" the entire
15 surface into the gas phase. Furthermore, note that some EAM
16 when used as free chemicals to embed analyte molecules as
17 described for the MALDI process will not work (i.e., they do
18 not promote molecular desorption, thus they are not suitable
19 matrix molecules).

20 As used herein, "probe element" or "sample presenting
21 device" refers to an element having the following properties:
22 it is inert (for example, typically stainless steel) and
23 active (probe elements with surfaces enhanced to contain EAM

1 and/or molecular capture devices).

2 As used herein, "MALDI" refers to Matrix-Assisted Laser
3 Desorption/Ionization.

4 As used herein, "TOF" stands for Time-of-Flight.

5 As used herein, "MS" refers to Mass Spectrometry.

6 As used herein, "MS/MS" refers to multiple sequential
7 mass spectrometry.

8 As used herein "MALDI-TOF MS" refers to Matrix-assisted
9 laser desorption/ionization time-of-flight mass spectrometry.

10 As used herein, "ESI" is an abbreviation for
11 electrospray ionization.

12 As used herein, "chemical bonds" is used simply as an
13 attempt to distinguish a rational, deliberate, and
14 knowledgeable manipulation of known classes of chemical
15 interactions from the poorly defined kind of general
16 adherence observed when one chemical substance (e.g., matrix)
17 is placed on another substance (e.g., an inert probe element
18 surface). Types of defined chemical bonds include
19 electrostatic or ionic (+/-) bonds (e.g., between a
20 positively and negatively charged groups on a protein
21 surface), covalent bonds (very strong or "permanent" bonds
22 resulting from true electron sharing), coordinate covalent
23 bonds (e.g., between electron donor groups in proteins and

1 transition metal ions such as copper or iron), and
2 hydrophobic interactions (such as between two noncharged
3 groups), weak dipole and London force or induced dipole
4 interactions.

5 As used herein, "electron donor groups" refers to the
6 case of biochemistry, where atoms in biomolecules (e.g, N, S,
7 O) "donate" or share electrons with electron poor groups
8 (e.g., Cu ions and other transition metal ions).

9 As used herein, the term "biopolymer markers indicative
10 or predictive of a disease state" is interpreted to mean that
11 a biopolymer marker which is strongly present in a normal
12 individual, but is down-regulated in disease is predictive of
13 said disease; while alternatively, a biopolymer marker which
14 is strongly present in a disease state, but is down-regulated
15 in normal individuals, is indicative of said disease state.
16 Biopolymer markers which are present in both disease and
17 normal states are indicative/predictive based upon their
18 relative strengths in disease vs. normal, along with the
19 observation regarding when their signal strengthens/weakens
20 relative to disease manifestation or progression.

21 As used herein, the term "disease state assessment" is
22 interpreted to mean quantitative or qualitative determination
23 of the presence/absence of the disease, with or without an

1 ability to determine severity, rapidity of onset, or
2 resolution of the disease state, e.g. a return to a normal
3 physiological state.

4 As used herein, the term "therapeutic target
5 recognition, development, and validation" refers to any
6 concept or method which enables an artisan to recognize,
7 develop, or validate the efficacy of a therapeutic moiety
8 which is effected in conjunction with a chemical or physical
9 interaction with one or more of the biopolymer markers of the
10 instant invention.

11 As used herein, the term "polypeptide" is interpreted to
12 mean a polymer composed of amino acid residues, related
13 naturally occurring structural variants, and synthetic non-
14 naturally occurring analogs thereof linked via peptide bonds,
15 related naturally occurring structural variants, and
16 synthetic non-naturally occurring analogs thereof. Synthetic
17 polypeptides can be synthesized, for example, using an
18 automated polypeptide synthesizer. The term "protein"
19 typically refers to large polypeptides. The term "peptide"
20 typically refers to short polypeptides. "Polypeptide(s)"
21 refers to any peptide or protein comprising two or more amino
22 acids joined to each other by peptide bonds or modified
23 peptide bonds. "Polypeptide(s)" refers to both short chains,

1 hybridize. The binding partner may itself be directly
2 detectable, for example, an antibody may be itself labeled
3 with a fluorescent molecule. The binding partner also may be
4 indirectly detectable, for example, a nucleic acid having a
5 complementary nucleotide sequence can be a part of a branched
6 DNA molecule that is in turn detectable through hybridization
7 with other labeled nucleic acid molecules. (See, e.g., P. D.
8 Fahrlander and A. Klausner, Bio/Technology (1988) 6:1165.)
9 Quantitation of the signal is achieved by, e.g.,
10 scintillation counting, densitometry, or flow cytometry.

11 As used herein, the term "antibody or antibodies"
12 includes polyclonal and monoclonal antibodies of any isotype
13 (IgA, IgG, IgE, IgD, IgM), or an antigen-binding portion
14 thereof, including but not limited to F(ab) and Fv fragments,
15 single chain antibodies, chimeric antibodies, humanized
16 antibodies, and a Fab expression library. "Antibody" refers
17 to a polypeptide ligand substantially encoded by an
18 immunoglobulin gene or immunoglobulin genes, or fragments
19 thereof, which specifically binds and recognizes an epitope
20 (e.g., an antigen). The recognized immunoglobulin --genes
21 include the kappa and lambda light chain constant region
22 genes, the alpha, gamma, delta, epsilon and mu heavy chain
23 constant region genes, and the myriad immunoglobulin variable

1 analyte in a sample of heterogeneous compounds. Thus, under
2 designated assay (e.g., immunoassay) conditions, the ligand
3 or receptor binds preferentially to a particular analyte and
4 does not bind in a significant amount to other compounds
5 present in the sample. For example, a polynucleotide
6 specifically binds under hybridization conditions to an
7 analyte polynucleotide comprising a complementary sequence;
8 an antibody specifically binds under immunoassay conditions
9 to an antigen analyte bearing an epitope against which the
10 antibody was raised; and an adsorbent specifically binds to
11 an analyte under proper elution conditions.

12 As used herein, the term "pharmaceutically effective
13 carrier" refers to any solid or liquid material which may be
14 used in creating formulations that further include active
15 ingredients of the instant invention, e.g. biopolymer markers
16 or therapeutics, for administration to a patient.

17 As used herein, the term "agent" is interpreted to mean
18 a chemical compound, a mixture of chemical compounds, a
19 sample of undetermined composition, a combinatorial small
20 molecule array, a biological macromolecule, a bacteriophage
21 peptide display library, a bacteriophage antibody (e.g.,
22 scFv) display library, a polysome peptide display library, or
23 an extract made from biological materials such as bacteria,

1 plants, fungi, or animal cells or tissues. Suitable
2 techniques involve selection of libraries of recombinant
3 antibodies in phage or similar vectors. See, Huse et al.
4 (1989) Science 246: 1275-1281; and Ward et al. (1989) Nature
5 341: 544-546. The protocol described by Huse is
6 rendered more efficient in combination with phage display
7 technology. See, e.g., Dower et al., WO 91/17271 and
8 McCafferty et al., WO 92/01047.

9 As used herein, the term "isolated" is interpreted to
10 mean altered "by the hand of man" from its natural state,
11 i.e., if it occurs in nature, it has been changed or removed
12 from its original environment, or both. For example, a
13 polynucleotide or a polypeptide naturally present in a living
14 organism is not "isolated," but the same polynucleotide or
15 polypeptide separated from the coexisting materials of its
16 natural state is "isolated", as the term is employed herein.

17 As used herein, the term "variant" is interpreted to
18 mean a polynucleotide or polypeptide that differs from a
19 reference polynucleotide or polypeptide respectively, but
20 retains essential properties. A typical variant of a
21 polynucleotide differs in nucleotide sequence from another,
22 reference polynucleotide. Changes in the nucleotide sequence
23 of the variant may or may not alter the amino acid sequence

1 immunologically reactive fragments, variants or moieties
2 thereof.

3 As used herein, the term "fragment" refers to the
4 products of the chemical, enzymatic, or physical breakdown of
5 an analyte. Fragments may be in a neutral or ionic state.

6 As used herein, the term "therapeutic avenues" is
7 interpreted to mean any agents, modalities, synthesized
8 compounds, etc., which interact with a biopolymer marker in
9 any manner that facilitates a therapeutic benefit, including
10 immunotherapeutic intervention, e.g. modalities such as
11 administration of an immunologically reactive moiety capable
12 of altering the course, progression and/or manifestation of
13 the disease, as a result of interfering with the disease
14 manifestation process, for example, at the early stages
15 focused upon by the identification of the disease, such as by
16 supplying a moiety capable of modifying the pathogenicity of
17 lymphocytes specific for the biopolymer marker or related
18 components.

19 As used herein, the term "interacting with a biopolymer
20 marker" includes any process by which a biopolymer marker may
21 physically or chemically relate with an organism,
22 particularly when this interaction results in the development
23 of therapeutic avenues or in modulation of the disease state.

1 ability to reduce the pathogenicity of key lymphocytes which
2 are specific for the native epitope. An analog is defined as
3 having structural similarity but not identity in peptide
4 sequencing able to be recognized by T-cells spontaneously
5 arising and targeting the endogeneous self epitope. A
6 critical function of this analog is an altered T-cell
7 activation which leads to T-cell anergy or death.

8 With the advent of mass spectrometric methods such as
9 MALDI and SELDI and ESI, researchers have begun to utilize a
10 tool that holds the promise of uncovering countless
11 biopolymers which result from translation, transcription and
12 post-translational transcription of proteins from the entire
13 genome.

14 Operating upon the principles of retentate
15 chromatography, SELDI MS involves the adsorption of proteins,
16 based upon their physico-chemical properties at a given pH
17 and salt concentration, followed by selectively desorbing
18 proteins from the surface by varying pH, salt, or organic
19 solvent concentration. After selective desorption, the
20 proteins retained on the SELDI surface, the "chip", can be
21 analyzed using the CIPHERGEN protein detection system, or an
22 equivalent thereof. Retentate chromatography is limited,
23 however, by the fact that if unfractionated body fluids, e.g.

1 blood, blood products, urine, saliva, cerebrospinal fluid,
2 lymph and the like, along with tissue samples, are applied
3 to the adsorbent surfaces, the biopolymers present in the
4 greatest abundance will compete for all the available binding
5 sites and thereby prevent or preclude less abundant
6 biopolymers from interacting with them, thereby reducing or
7 eliminating the diversity of biopolymers which are readily
8 ascertainable.

9 If a process could be devised for maximizing the
10 diversity of biopolymers discernable from a sample, the
11 ability of researchers to accurately determine the relevance
12 of such biopolymers with relation to one or more disease
13 states would be immeasurably enhanced.

14
15 SUMMARY OF THE INVENTION

16 The instant invention is characterized by the use of a
17 combination of preparatory steps, e.g. chromatography and 1-D
18 tricine polyacrylamide gel electrophoresis. Subsequent to
19 which the gel is stained, e.g. with Coomassie blue, silver or
20 rubidium. Next, bands are selected from the gels for further
21 study. Tryptic digestion of each band follows, concluding
22 with the extraction of tryptic peptides from the digest.
23 This extraction may be accomplished utilizing C18 ZIPTIPs, or

1 organic extract and dry technique followed by MALDI Qq TOF
2 (Maldi Quadrupole Quadrupole Time of Flight) processing.
3 Additional methodologies may include SELDI MS, 2-D gel
4 technology, MALDI MS/MS and time-of-flight detection
5 procedures to maximize the diversity of biopolymers which are
6 verifiable within a particular sample. The cohort of
7 biopolymers verified within a sample is then compared to
8 develop data indicating their presence, absence or relative
9 strength/concentration in disease vs normal controls, and
10 further studied to determine whether the up-regulation or
11 down-regulation of a single biopolymer or group of
12 biopolymers is indicative of a disease state or predictive of
13 the development of said disease state. Additionally,
14 biopolymers recognized as being indicative or predictive of a
15 disease state in accordance with the instant invention are
16 useful in therapeutic intervention, e.g. as therapeutic
17 modalities in their own right, in the course of therapeutic
18 target recognition, in the development and validation of
19 efficacious therapeutic modalities, e.g. when interrogating or
20 developing phage display libraries, and as ligands or
21 receptors for use in conjunction with therapeutic
22 intervention.

23 Although all manner of biomarkers related to all disease

1 conditions are deemed to be within the purview of the instant
2 invention and methodology, particular significance was given
3 to those markers and diseases associated with the complement
4 system, cognitive diseases, e.g. Alzheimer's disease and
5 Syndrome X and diseases related thereto.

6 The complement system is an important part of non-clonal
7 or innate immunity that collaborates with acquired immunity
8 to destroy invading pathogens and to facilitate the clearance
9 of immune complexes from the system. This system is the
10 major effector of the humoral branch of the immune system,
11 consisting of nearly 30 serum and membrane proteins. The
12 proteins and glycoproteins composing the complement system
13 are synthesized largely by liver hepatocytes. Activation of
14 the complement system involves a sequential enzyme cascade in
15 which the proenzyme product of one step becomes the enzyme
16 catalyst of the next step. Complement activation can occur
17 via two pathways: the classical and the alternative. The
18 classical pathway is commonly initiated by the formation of
19 soluble antigen-antibody complexes or by the binding of
20 antibody to antigen on a suitable target, such as a bacterial
21 cell. The alternative pathway is generally initiated by
22 various cell-surface constituents that are foreign to the
23 host. Each complement component is designated by numerals

1 (C1-C9), by letter symbols, or by trivial names. After a
2 component is activated, the peptide fragments are denoted by
3 small letters. The complement fragments interact with one
4 another to form functional complexes. Ultimately, foreign
5 cells are destroyed through the process of a membrane-attack
6 complex mediated lysis.

7 The C4 component of the complement system is involved in
8 the classical activation pathway. It is a glycoprotein
9 containing three polypeptide chains (α , β , and γ). C4 is a
10 substrate of component C1s and is activated when C1s
11 hydrolyzes a small fragment (C4a) from the amino terminus of
12 the α chain, exposing a binding site on the larger fragment
13 (C4b).

14 The native C3 component consists of two polypeptide
15 chains, α and β . As a serum protein, C3 is involved in the
16 alternative pathway. Serum C3, which contains an unstable
17 thioester bond, is subject to slow spontaneous hydrolysis
18 into C3a and C3b. The C3f component is involved in the
19 regulation required of the complement system which confines
20 the reaction to designated targets. During the regulation
21 process, C3b is cleaved into two parts: C3bi and C3f. C3bi
22 is a membrane-bound intermediate wherein C3f is a free
23 diffusible (soluble) component.

1 Complement components have been implicated in the
2 pathogenesis of several disease conditions. C3 deficiencies
3 have the most severe clinical manifestations, such as
4 recurrent bacterial infections and immune-complex diseases,
5 reflecting the central role of C3. The rapid profusion of
6 C3f moieties and resultant "accidental" lysis of normal cells
7 mediated thereby gives rise to a host of auto-immune
8 reactions. The ability to understand and control these
9 mechanisms, along with their attendant consequences, will
10 enable practitioners to develop both diagnostic and
11 therapeutic avenues by which to thwart these maladies.

12 In the course of defining a plurality of disease
13 specific marker sequences, special significance was given to
14 markers which were evidentiary of a particular disease state
15 or with conditions associated with Syndrome-X. Syndrome-X is
16 a multifaceted syndrome, which occurs frequently in the
17 general population. A large segment of the adult population
18 of industrialized countries develops this metabolic syndrome,
19 produced by genetic, hormonal and lifestyle factors such as
20 obesity, physical inactivity and certain nutrient excesses.
21 This disease is characterized by the clustering of insulin
22 resistance and hyperinsulinemia, and is often associated with
23 dyslipidemia (atherogenic plasma lipid profile), essential

1 hypertension, abdominal (visceral) obesity, glucose
2 intolerance or noninsulin-dependent diabetes mellitus and an
3 increased risk of cardiovascular events. Abnormalities of
4 blood coagulation (higher plasminogen activator inhibitor
5 type I and fibrinogen levels), hyperuricemia and
6 microalbuminuria have also been found in metabolic syndrome-
7 X.

8 The instant inventors view the Syndrome X continuum in
9 its cardiovascular light, while acknowledging its important
10 metabolic component. The first stage of Syndrome X consists
11 of insulin resistance, abnormal blood lipids (cholesterol,
12 triglycerides and free fatty acids), obesity, and high blood
13 pressure (hypertension). Any one of these four first stage
14 conditions signals the start of Syndrome X.

15 Each first stage Syndrome X condition risks leading to
16 another. For example, increased insulin production is
17 associated with high blood fat levels, high blood pressure,
18 and obesity. Furthermore, the effects of the first stage
19 conditions are additive; an increase in the number of
20 conditions causes an increase in the risk of developing more
21 serious diseases on the Syndrome X continuum.

22 A patient who begins the Syndrome X continuum risks
23 spiraling into a maze of increasingly deadly diseases. The

1 next stages of the Syndrome X continuum lead to overt
2 diabetes, kidney failure, and heart failure, with the
3 possibility of stroke and heart attack at any time. Syndrome
4 X is a dangerous continuum, and preventative medicine is the
5 best defense. Diseases are currently most easily diagnosed
6 in their later stages, but controlling them at a late stage
7 is extremely difficult. Disease prevention is much more
8 effective at an earlier stage.

9 In a further contemplated embodiment of the invention,
10 samples may be taken from a patient at one point in time, as
11 a single sample or as multiple samples, or at different
12 points in time such that analysis is carried out on multiple
13 samples for ongoing analysis. Typically, a first sample is
14 taken from a patient upon presentation with possible symptoms
15 of a disease and analyzed according to the invention.
16 Subsequently, some period of time after presentation, for
17 example, about 3 - 6 months after the first presentation, a
18 second sample is taken and analyzed according to the
19 invention. The data can be used, by way of example, to
20 diagnose or monitor a disease state, determine risk
21 assessment, identify therapeutic avenues, or determine the
22 therapeutic value of an agent such as a pharmaceutical.

23 Subsequent to the isolation of particular disease state

1 marker sequences as taught by the instant invention, the
2 promulgation of various forms of risk assessment tests are
3 contemplated which will allow physicians to identify
4 asymptomatic patients before they suffer an irreversible
5 event such as diabetes, kidney failure, and heart failure,
6 and enable effective disease management and preventative
7 medicine. Additionally, the specific diagnostic tests which
8 evolve from this methodology provide a tool for rapidly and
9 accurately diagnosing acute Syndrome X events such as heart
10 attack and stroke, and facilitate treatment.

11 More particularly, biopolymer markers elucidated via
12 methodologies of the instant invention find utility related
13 to broad areas of disease therapeutics. Such therapeutic
14 avenues include, but are not limited to:

15 1) utilization and recognition of said biopolymer
16 markers, variants or moieties thereof as direct therapeutic
17 modalities, either alone or in conjunction with an effective
18 amount of a pharmaceutically effective carrier;

19 2) validation of therapeutic modalities or disease
20 preventative agents as a function of biopolymer marker
21 presence or concentration;

22 3) treatment or prevention of a disease state by
23 formation of disease intervention modalities; e.g. formation

1 of biopolymer/ligand conjugates which intervene at receptor
2 sites to prevent, delay or reverse a disease process;

3 4) use of biopolymer markers or moieties thereof as a
4 means of elucidating therapeutically viable agents, e.g. from
5 a bacteriophage peptide display library, a bacteriophage
6 antibody library or the like;

7 5) instigation of a therapeutic immunological
8 response; and

9 6) synthesis of molecular structures related to said
10 biopolymer markers, moieties or variants thereof which are
11 constructed and arranged to therapeutically intervene in the
12 disease process.

13 A process for identifying or developing therapeutic
14 avenues related to a disease state utilizing any of the above
15 examples may follow results obtained from conducting an
16 analysis inclusive of interacting with a biopolymer including
17 the sequence of the particular disease specific marker or at
18 least one analyte thereof of the present invention. Such
19 treatment or prevention of a disease state by formation of
20 disease intervention modalities may be by the formation of
21 biopolymer/ligand conjugates which intervene at receptor
22 sites to prevent, delay, or reverse a disease process. In
23 addition, a means of elucidating therapeutically viable

1 agents may include the use of a bacteriophage peptide display
2 library or a bacteriophage antibody library. The therapeutic
3 avenues may regulate the presence or absence of the
4 biopolymer including the sequence of the particular disease
5 specific marker or at least one analyte thereof in the
6 present invention.

7 Accordingly, it is an objective of the instant invention
8 to define a disease specific biopolymer marker sequence which
9 is useful in evidencing and categorizing at least one
10 particular disease state.

11 It is an additional objective of the instant invention
12 to develop methods and means of disease therapy, including
13 but not limited to:

14 1) utilization and recognition of said biopolymer
15 markers, variants or moieties thereof as direct therapeutic
16 modalities, either alone or in conjunction with an effective
17 amount of a pharmaceutically effective carrier;

18 2) validation of therapeutic modalities or disease
19 preventative agents as a function of biopolymer marker
20 presence or concentration;

21 3) treatment or prevention of a disease state by
22 formation of disease intervention modalities; e.g. formation
23 of biopolymer/ligand conjugates which intervene at receptor

1 sites to prevent, delay or reverse a disease process;

2 4) use of biopolymer markers or moieties thereof as a
 3 means of elucidating therapeutically viable agents, e.g. from
 4 a bacteriophage peptide display library, a bacteriophage
 5 antibody library or the like;

6 5) instigation of a therapeutic immunological
 7 response; and

8 6) synthesis of molecular structures related to said
 9 biopolymer markers, moieties or variants thereof which are
 10 constructed and arranged to therapeutically intervene in the
 11 disease process, e.g. by directly determining the three-
 12 dimensional structure of said biopolymer marker directly from
 13 an amino acid sequence thereof.

14 It is another objective of the instant invention to
 15 evaluate samples containing a plurality of biopolymers for
 16 the presence of disease specific biopolymer marker sequences
 17 (disease specific markers) which evidence a link to at least
 18 one specific disease state.

19 It is a further objective of the instant invention to
 20 elucidate essentially all biopolymeric markers, moieties or
 21 variants thereof contained within said samples, whereby
 22 particularly significant moieties may be identified.

23 It is a further objective of the instant invention

1 provide at least one purified antibody which is specific to
2 said disease specific marker sequence.

3 It is yet another objective of the instant invention to
4 teach a monoclonal antibody which is specific to said disease
5 specific marker sequence.

6 It is a still further objective of the invention to
7 teach polyclonal antibodies raised against said disease
8 specific marker.

9 It is yet an additional objective of the instant
10 invention to teach a diagnostic kit for determining the
11 presence, concentration, or relative strength/concentration
12 of said disease specific marker.

13 It is a still further objective of the instant invention
14 to teach methods for characterizing disease state based upon
15 the identification of said disease specific marker.

16 Other objects and advantages of this invention will
17 become apparent from the following description taken in
18 conjunction with the accompanying drawings wherein are set
19 forth, by way of illustration and example, certain
20 embodiments of this invention. The drawings constitute a
21 part of this specification and include exemplary embodiments
22 of the present invention and illustrate various objects and
23 features thereof.

1

2 BRIEF DESCRIPTION OF THE FIGURES

3 Figure 1 is a photograph of a tricine gel DEAE 3 (Elution)
4 comparing Alzheimers disease versus Age Matched Control.

5

6 DETAILED DESCRIPTION OF THE INVENTION

7 In earlier work, for example in U.S. Patent application
8 09/846330 filed April 30, 2000, the contents of which is
9 herein incorporated by reference, raw sera was obtained and
10 mixed with formic acid and extracted the peptides with C18
11 reversed phase ZIPTIPs.

12 In the instantly disclosed invention, we deal with
13 proteins generally having a molecular weight of about 20 kD
14 or more. In general, proteins of greater than 20 kD can
15 reliably be fragmented by trypsin or other enzymes. The
16 instant technology incorporates sufficient sensitivity to
17 deal with even the low production of peptides from proteins
18 less than 20 kD clipped from gel.

19 Proteins differ from peptides in that they cannot be
20 effectively resolved by time of flight MS and they are too
21 large (>3kD) to be effectively fragmented by collision with
22 gases. The most commonly used solution to these problems is
23 to resolve the proteins by polyacrylamide gel electrophoresis

FOCET 662660

1 followed by staining with silver, or coomasie brilliant blue
2 or rubidium dyes or counter staining with Zinc-SDS complexes.
3 Once the proteins have been resolved and visualized with
4 stains the proteins that differ between disease states can
5 then be excised from the gel and the protein purified in the
6 1-D gel band or 2-D gel spot can be cleaved into fragments
7 less than 3 kD by proteolytic enzymes. Once protein has been
8 resolved by gel and cleaved by enzymes, the protein is
9 considered in the form of peptides and therefore can be dealt
10 with as per earlier work (09/846330). The peptide is either
11 collected and purified with C18 reversed phase chromatography
12 or by some other form of chromatography prior to reversed
13 phase separation. The peptide can also be collected in
14 ammonium carbonate buffer that is subsequently evolved by
15 reaction with acid or by removal in organic solvents.

16 Once the peptides are collected they can be sequenced,
17 e.g. with a MALDI-Qq-TOF but also with a TOF-TOF, and
18 ESI-Q-TOF or an ION-TRAP. Other types of MS analysis which
19 may be employed are SELDI MS and MS/MS. The peptides are
20 fragments of the original protein. The peptides are
21 sequenced by fragmentation to produce a spectrum composed of
22 the parts of the peptide. The peptide fragments can be
23 produced by a strong ionization energy with a laser,

1 human blood-borne protein, is thus identified. It is possible
2 to identify plural targets in this fashion.

3

4 Following are exemplary, but non-limiting examples of
5 preparatory protocols useful in the process of the instant
6 invention.

7

8 Preparatory Protocols:

9 Any of these protocols may be selected from a column
10 flow-through stream, a column elution stream, or a column
11 scrub stream.

12 Hi Q is a strong anion exchanger made of methyl acrylate
13 co-polymer with the functional group: $-N^+(CH_3)_2$;

14 Hi S is a strong cation exchanger made of methyl acrylate
15 co-polymer with the functional group: $-SO_3^-$;

16 DEAE is diethylaminoethyl which is a weak cation exchanger
17 made of methyl acrylate co-polymer with the functional group
18 $-N^+(C_2H_5)_2$;

19 PS is phenyl sepharose;

20 BS is butyl sepharose.

21 Note that the supports, i.e. methyl acrylate and
22 sepharose are different, but non-limiting examples, as the
23 same functional group on different supports will function,

1 albeit possibly with different effects.

2 DEAE Column Protocol:

- 3 1)Cast 200 µl of 50% slurry;
4 2)Equilibrate column in 5 bed volumes of 50 mM
5 tricine pH 8.8 (binding buffer);
6 3)Dissolve 25 µl of sera in 475 µl of binding buffer;
7 4)Wash column in 5 bed volumes of binding buffer;
8 5)Elute column in 120 µl of 0.4 M Phosphate buffer
9 (PB) pH 6.1;
10 6)Elute column in 120 µl of 50 mM citrate buffer
11 pH 4.2;
12 7)Scrub column with 120 µl sequentially with each
13 of 0.1% triton, 1.0% triton and 2% SDS in
14 62.5 mM Tris pH 6.8.

16 Butyl Sepharose Column Protocol:

- 17 1)Cast 150 µl bed volume column;
18 2)Equilibrate column in 5 bed volumes of 1.7 M
19 $(\text{NH}_4)_2\text{SO}_4$ in 50 mM PB pH 7.0 (binding buffer);
20 3)Dissolve 35 µl of sera in 465 µl of binding buffer
21 and apply;
22 4)Wash column in 5 bed volumes of binding buffer;
23 5)Elute column in 120 µl of 0.4 M $(\text{NH}_4)_2\text{SO}_4$ in

- 1 50 mM PB pH 7.0;
2 6) Elute column in 120 µl of 50 mM PB pH 7.0;
3 7) Scrub column with 120 µl sequentially with each
4 of 0.1% triton, 1.0% triton and 2% SDS in
5 62.5 mM Tris pH 6.8.

6
7
8 Phenyl Sepharose Column Protocol:

- 9 1) Cast 150 µl bed volume column;
10 2) Equilibrate column in 5 bed volumes of
11 1.7 M $(\text{NH}_4)_2\text{SO}_4$ in 50 mM PB pH 7.0 (binding buffer);
12 3) Dissolve 35 µl of sera in 465 µl of binding
13 buffer and apply;
14 4) Wash column in 5 bed volumes of binding buffer;
15 5) Elute column in 120 µl of 0.2 M $(\text{NH}_4)_2\text{SO}_4$ in
16 50 mM PB pH 7.0;
17 6) Elute column in 120 µl of 50 mM PB pH 7.0;
18 7) Scrub column with 120 µl sequentially with each
19 of 0.1% triton, 1.0% triton and 2% SDS in
20 62.5 mM Tris pH 6.8.

21
22 HiO Anion Exchange Mini Column Protocol:

- 23 1) Dilute sera in sample/running buffer;

- 1 2)Add HiQ resin to column and remove any air bubbles;
2 3)Add ultrafiltered (UF) water to aid in column
3 packing;
4 4)Add sample/running buffer to equilibrate column;
5 5)Add diluted sera;
6 6)Collect all the flow-through fraction in Eppendorf
7 tubes until level is at resin;
8 7)Add sample/running buffer to wash column;
9 8)Add elution buffer and collect elution in Eppendorf
10 tubes.

11
12 HiS Cation Exchange Mini Column Protocol:

- 13 1)Dilute sera in sample/running buffer;
14 2)Add HiS resin to column and remove any air bubbles;
15 3)Add UF water to aid in column packing;
16 4)Add sample/running buffer to equilibrate column for
17 sample loading;
18 5)Add diluted sera to column;
19 6)Collect all flow through fractions in Eppendorf
20 tubes until level is at resin;
21 7)Add sample/running buffer to wash column;
22 8)Add elution buffer and collect elution in Eppendorf
23 tubes.

0991664660
FOEET" 664660

1 Illustrative of the various buffering compositions
2 useful in this technique are:
3 Sample/Running buffers: including but not limited to
4 Bicine buffers of various molarities, pH's, NaCl content,
5 Bis-Tris buffers of various molarities, pH's, NaCl
6 content, Diethanolamine of various molarities, pH's, NaCl
7 content, Diethylamine of various molarities, pH's, NaCl
8 content, Imidazole of various molarities, pH's, NaCl
9 content, Tricine of various molarities, pH's, NaCl
10 content, Triethanolamine of various molarities, pH's, NaCl
11 content, Tris of various molarities, pH's, NaCl content.
12 Elution Buffer: Acetic acid of various molarities, pH's,
13 NaCl content, Citric acid of various molarities, pH's,
14 NaCl content, HEPES of various molarities, pH's, NaCl
15 content, MES of various molarities, pH's, NaCl content,
16 MOPS of various molarities, pH's, NaCl content, PIPES of
17 various molarities, pH's, NaCl content, Lactic acid of
18 various molarities, pH's, NaCl content, Phosphate of
19 various molarities, pH's, NaCl content, Tricine of various
20 molarities, pH's, NaCl content.

21 Following tryptic digestion, additional processing
22 may be carried out, for example:
23 Utilizing a type of micro-chromatographic column called a

1 C18- ZIPTIP available from the Millipore company, the
2 following preparatory steps were conducted.

- 3 1. Dilute sera in sample buffer
- 4 2. Aspirate and dispense ZIPTIP in 50% Acetonitrile
- 5 3. Aspirate and dispense ZIPTIP in Equilibration solution
- 6 4. Aspirate and dispense in serum sample
- 7 5. Aspirate and dispense ZIPTIP in Wash solution
- 8 6. Aspirate and dispense ZIPTIP in Elution Solution

9 Illustrative of the various buffering compositions
10 useful in the present invention are:

11 Sample Buffers (various low pH's): Hydrochloric acid
12 (HCl), Formic acid, Trifluoroacetic acid (TFA),

13 Equilibration Buffers (various low pH's): HCl, Formic
14 acid, TFA;

15 Wash Buffers (various low pH's): HCl, Formic acid, TFA;

16 Elution Solutions (various low pH's and % Solvents):

17 HCl, Formic acid, TFA;

18 Solvents: Ethanol, Methanol, Acetonitrile.

19 Spotting was then performed, for example upon a Gold Chip
20 in the following manner:

- 21 1. Spot 2 ul of sample onto each spot
- 22 2. Let sample partially dry

23 As a result of these procedures, the disease specific

Dwain

1 markers plasma protease (C1) inhibitor having a molecular
2 weight of about 1826 daltons and a sequence of
3 (K)GVTSVSQIFHSPDLAIR(D), molecular weight of about 1560
4 daltons and a sequence of (K)KYPVAHFIDQTLK(A), molecular
5 weight of about 1186 and a sequence of (K)FQPTLLTLPR(I) (see
6 Fig. 1, Band 2) related to Alzheimers disease were found.

7 Figure 1 is a photograph of a gel which is indicative
8 of the presence/absence of the marker in disease vs. control
9 and, in cases where the marker is always present, the
10 relative strength, e.g. the up or down regulation of the
11 marker relative to categorization of disease state is
12 deduced.

13 A method for evidencing and categorizing at least one
14 disease state is disclosed. The steps taken include
15 obtaining a sample from a patient, preferably human, and
16 conducting MS analysis on the sample. As a result, at least
17 one biopolymer marker sequence or analyte thereof is
18 isolated from the sample which undergoes evidencing and
19 categorizing and is compared to the biopolymer marker
20 sequence as disclosed in the present invention. The step of
21 evidencing and categorizing is particularly directed to
22 biopolymer markers or analytes thereof linked to at least
23 one risk of disease development of the patient or related to

1 the existence of a particular disease state.

2 In addition, various kits are contemplated for use by
3 the present invention. One such kit provides for
4 determining the presence of the disease specific biopolymer
5 marker. At least one biochemical material is incorporated
6 which is capable of specifically binding with a biomolecule
7 which includes at least the disease specific biopolymer
8 marker or analyte thereof, and a means for determining
9 binding between the biochemical material and the
10 biomolecule. The biochemical material for any of the
11 contemplated kits, by way of example an antibody or at least
12 one monoclonal antibody specific therefore, or biomolecule
13 may be immobilized on a solid support and include at least
14 one labeled biochemical material which is preferably an
15 antibody. The sample utilized for any of the kits may be a
16 fractionated or unfractionated body fluid or a tissue
17 sample. Non-limiting examples of such fluids are blood,
18 blood products, urine, saliva, cerebrospinal fluid, and
19 lymph.

20 Further contemplated is a kit for diagnosing,
21 determining risk-assessment, and identifying therapeutic
22 avenues related to a disease state. This kit includes at
23 least one biochemical material which is capable of

09991799 "113301
1 specifically binding with a biomolecule which includes at
2 least one biopolymer marker including the sequence of the
3 particular disease specific biopolymer marker or an
4 analyte thereof related to the disease state. Also
5 included is a means for determining binding between the
6 biochemical material and the biomolecule, whereby at least
7 one analysis to determine a presence of a marker, analyte
8 thereof, or a biochemical material specific thereto, is
9 carried out on a sample. As previously described,
10 analysis may be carried out on a single sample or multiple
11 samples.

12 In accordance with various stated objectives of the
13 invention, the skilled artisan, in possession of the
14 specific disease specific marker as instantly disclosed,
15 would readily carry out known techniques in order to raise
16 purified biochemical materials, e.g. monoclonal and/or
17 polyclonal antibodies, which are useful in the production of
18 methods and devices useful as point-of-care rapid assay
19 diagnostic or risk assessment devices as are known in the
20 art.

21 The specific disease markers which are analyzed
22 according to the method of the invention are released into
23 the circulation and may be present in the blood or in any

1 blood product, for example plasma, serum, cytolyzed blood,
2 e.g. by treatment with hypotonic buffer or detergents and
3 dilutions and preparations thereof, and other body fluids,
4 e.g. CSF, saliva, urine, lymph, and the like. The
5 presence of each marker is determined using antibodies
6 specific for each of the markers and detecting specific
7 binding of each antibody to its respective marker. Any
8 suitable direct or indirect assay method may be used to
9 determine the level of each of the specific markers
10 measured according to the invention. The assays may be
11 competitive assays, sandwich assays, and the label may be
12 selected from the group of well-known labels such as
13 radioimmunoassay, fluorescent or chemiluminescence
14 immunoassay, or immunoPCR technology. Extensive discussion
15 of the known immunoassay techniques is not required here
16 since these are known to those of skilled in the art. See
17 Takahashi et al. (Clin Chem 1999;45(8):1307) for a
18 detailed example of an assay.

19 A monoclonal antibody specific against the disease
20 marker sequence isolated by the present invention may be
21 produced, for example, by the polyethylene glycol (PEG)
22 mediated cell fusion method, in a manner well-known in the
23 art.

09991799.12301
FOEET 667660

1 Traditionally, monoclonal antibodies have been made
2 according to fundamental principles laid down by Kohler
3 and Milstein. Mice are immunized with antigens, with or
4 without, adjuvants. The splenocytes are harvested from
5 the spleen for fusion with immortalized hybridoma
6 partners. These are seeded into microtiter plates where
7 they can secrete antibodies into the supernatant that is
8 used for cell culture. To select from the hybridomas that
9 have been plated for the ones that produce antibodies of
10 interest, the hybridoma supernatants are usually tested
11 for antibody binding to antigens in an ELISA (enzyme
12 linked immunosorbent assay) assay. The idea is that the
13 wells that contain the hybridoma of interest will contain
14 antibodies that will bind most avidly to the test antigen,
15 usually the immunizing antigen. These wells are then
16 subcloned in limiting dilution fashion to produce
17 monoclonal hybridomas. The selection for the clones of
18 interest is repeated using an ELISA assay to test for
19 antibody binding. Therefore, the principle that has been
20 propagated is that in the production of monoclonal
21 antibodies the hybridomas that produce the most avidly
22 binding antibodies are the ones that are selected from
23 among all the hybridomas that were initially produced.

1 That is to say, the preferred antibody is the one with
2 highest affinity for the antigen of interest.

3 There have been many modifications of this procedure
4 such as using whole cells for immunization. In this
5 method, instead of using purified antigens, entire cells
6 are used for immunization. Another modification is the
7 use of cellular ELISA for screening. In this method
8 instead of using purified antigens as the target in the
9 ELISA, fixed cells are used. In addition to ELISA tests,
10 complement mediated cytotoxicity assays have also been
11 used in the screening process. However, antibody-binding
12 assays were used in conjunction with cytotoxicity tests.
13 Thus, despite many modifications, the process of producing
14 monoclonal antibodies relies on antibody binding to the
15 test antigen as an endpoint.

16 The purified monoclonal antibody is utilized for
17 immunochemical studies.

18 Polyclonal antibody production and purification
19 utilizing one or more animal hosts in a manner well-known
20 in the art can be performed by a skilled artisan.

21 Another objective of the present invention is to
22 provide reagents for use in diagnostic assays for the
23 detection of the particularly isolated disease specific

1 marker sequences of the present invention.

2 In one mode of this embodiment, the marker sequences
3 of the present invention may be used as antigens in
4 immunoassays for the detection of those individuals
5 suffering from the disease known to be evidenced by said
6 marker sequence. Such assays may include but are not
7 limited to: radioimmunoassay, enzyme-linked immunosorbent
8 assay (ELISA), "sandwich" assays, precipitin reactions,
9 gel diffusion immunodiffusion assay, agglutination assay,
10 fluorescent immunoassays, protein A or G immunoassays and
11 immunoelectrophoresis assays.

12 According to the present invention, monoclonal or
13 polyclonal antibodies produced against the disease
14 specific marker sequence of the instant invention are
15 useful in an immunoassay on samples of blood or blood
16 products such as serum, plasma or the like, cerebrospinal
17 fluid or other body fluid, e.g. saliva, urine, lymph, and
18 the like, to diagnose patients with the characteristic
19 disease state linked to said marker sequence. The
20 antibodies can be used in any type of immunoassay. This
21 includes both the two-site sandwich assay and the single
22 site immunoassay of the non-competitive type, as well as
23 in traditional competitive binding assays.

1 a tool for monitoring the progress of a patient during a
2 therapeutic treatment, so as to determine the usefulness
3 of a novel therapeutic agent.

4 All patents and publications mentioned in this
5 specification are indicative of the levels of those
6 skilled in the art to which the invention pertains. All
7 patents and publications are herein incorporated by
8 reference to the same extent as if each individual
9 publication was specifically and individually indicated to
10 be incorporated by reference.

11 It is to be understood that while a certain form of
12 the invention is illustrated, it is not to be limited to
13 the specific form or arrangement herein described and
14 shown. It will be apparent to those skilled in the art
15 that various changes may be made without departing from
16 the scope of the invention and the invention is not to be
17 considered limited to what is shown and described in the
18 specification and drawings/figures.

19 One skilled in the art will readily appreciate that
20 the present invention is well adapted to carry out the
21 objectives and obtain the ends and advantages mentioned,
22 as well as those inherent therein. The oligonucleotides,
23 peptides, polypeptides, biologically related compounds,

